Kills Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds; including those listed on the back panel.

Contains *TRIMEC® Herbicide to Control the Toughest Weeds

1 Bag Covers 15.000 Sg. Ft.

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid	0.55%
(+)-(R)-2-(2 methyl-4-chlorophenoxy) propionic acid	0.12%
Dicamba: 3,6-dichloro-o-anisic acid	0.05%
OTHER INGREDIENTS:	

EPA Reg. No. 2217-643-7001

EPA EST. NO. 69759-OR-004 (Lot begins BAN)

THIS PRODUCT CONTAINS:

11.0 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton. 2.4 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy) propionc acid equivalent per ton. 1.0 lb. 3,6-dichloro-o-anisic acid equivalent per ton. Isomer Specific by AOAC Methods.

KEEP OUT OF REACH OF CHILDREN CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material.

First Aid	
If on skin or on clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice
If inhaled:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

GUARANTEED ANALYSIS:

10.00/	TOTAL NUTBOOFN (NI)
	TOTAL NITROGEN (N)
	16.0% Ammoniacal Nitro
6.0%	AVAILABLE PHOSPHATE (P.
8.0%	
	16.0% Combined Sulfur (S
	IRON (Fe)
	MANGANESE (Mn)
hosphate, Ammonium Sulfate, Muriate of Potash, Iron Oxysulfate	Derived from Monoammoniur
	and Manganese Oxysulfate.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.html



NOTICE:This product can expose you to Cadmium (Ca), which is known to the State of California to cause cancer, birth defects or other reproductive harm. For more information go to www.P65Warnings.ca.gov.

SPREADER SETTINGS:

Apply 50 lbs. (for 15,000 sq. ft. coverage per bag)	SGN 150
BEST® Pro	4B
Spyker / BEST® Models 34, 44, 64	3.7
Earthway	12
Lesco	13
Scotts R-8A (Cone 8)	J½
1 1 40=0 (1)	

As a guide, set spreader at the setting indicated by the above chart before applying. Remember the settings provided are only a guide and depend on the age, condition and speed of operation of your spreader. Check spreader calibration often and make adjustment as needed. Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter product beyond lawn edge and onto concrete.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

• Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. • Do not apply this product to dichondra or carpetgrass. Do not apply this product to 'Floratam' St. Augustinegrass in Florida. • Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning to prevent over-application. • Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result. • Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or

street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

Turf Supreme® 16-6-8 Plus Trimec® controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns as listed. In addition, a 16-6-8 fertilizer provides plant nutrients or a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days.

The maximum broadcast application rate is 5.0 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.0 pounds of product per 1,000 sq. ft.

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

	Calibration Rate, Application Rate, Treate		Treated Area for th	is bag size, Sq. Ft.
Pounds	I	Pounds/1000 Sq. Ft.	Cool Season Grass	St. Augustinegrass
50	1/300	3.33	15,000	

WEEDS CONTROLLED

	WEEDO CONTROLLED	
Bindweed, field	Cocklebur	Prostrate knotweed
Bird vetch	Common mullein	Prostrate spurge
Bitter wintercress	Creeping buttercup	Puncturevine
Bittercress, hairy	Curly dock	Purslane, common
Black medic	Daisy, English	Red sorrel
Black mustard	Dandelion, common	Redroot pigweed
Blackseed plantain	Dichondra	Redstem filaree
Blessed thistle	Dollarweed	Rough cinquefoil
Bloodflower milkweed	Fiddleneck	Rough fleabane
Bracted plantain	Goldenrod	Russian pigweed
Bristly oxtongue	Ground ivy	Russian thistle
Broadleaf dock	Henbit	Scarlet pimpernel
Broadleaf plantain	Hoary cress	Scotch thistle
Buckhorn plantain	Hoary plantain	Shepard's purse
Bulbous buttercup	Kochia	Smooth dock
Bull thistle	Knotweed, prostrate	Smooth pigweed
Burdock	Lambsquarters	Sowthistle, annual
Burning nettle	Mallow, common	Speedwell, commor
Burclover	Morningglory, tall	Spotted spurge
Butonweed, common	Mustard, tumble	Tall Nettle
Canada thistle	Narrowleaf plantain	Tumble Pigweed
Carpetweed	Narrowleaf vetch	Venice mallow
Chickweed, common	Pearlwort	Wavyleaf, thistle
Chickweed, mouseear	Poison ivy	White mustard
Chicory	Poison oak	Wild carrot
Clover, crimson	Prickly lettuce	Wolly Plantain
Clover, white	Prostrate pigweed	Yarrow, common

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANT-ABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

*TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

Net Weight: 50 LBS. (22.68 Kg)



Distributed By:
J. R. SIMPLOT COMPANY
PROFESSIONAL PRODUCTS
P.O. Box 198 • CA 95330
www.bestfertilizer.com