



HERBICIDE/DESICCANT



GROWING POTATOES IS TOUGH, KILLING THEM CAN BE JUST AS HARD

You have a lot invested in your potato crop. From providing the optimal fertility and cultural practices to protecting from yield robbing insects, weeds and diseases, it can add up to a big investment. When you are finally ready for harvest to capture the results of all that work, it is important to have the right program to desiccate vines, kill any weeds, and prepare the crop for a successful harvest.

Innvictis Crop Care® has just made that task a little easier. The introduction of Terivate™ and Strevin™ 2EC, along with our new Inntero™ adjuvant technology, makes building an effective vine desiccation program straightforward.

HERBICIDE GROUP: 14



Strevin™ 2EC is a low use rate broadleaf contact herbicide. Strevin 2EC quickly controls many weeds like pigweed, lambsquarters, kochia and Russian thistle. Strevin 2EC is an excellent means of desiccating crops to aid in harvest operations.

Active Ingredient	APPLICATION
Carfentrazone	
Formulation	Ground
2.0 lbs ai/gal	20 gpa
Rate	Air
3.2 oz/A alone	5-10 gpa
Tank Mix	REI - 12 hrs
2.0-5.58 oz/A	

HERBICIDE GROUP: 22



Terivate™ is a preharvest, contact herbicide used to control late-season weeds while desiccating existing crop foliage. This combination allows for improved harvestability and grain quality.

Active Ingredient	APPLICATION
Diquat-dibromide	
Formulation	Ground: 20 gpa
2.0 lbs ai/gal	Air: 5 gpa
Rate	REI - 24 hrs
1-2 pts/A	

USE RECOMMENDATIONS

When potato crop begins to senesce, make desiccant application providing thorough coverage to all foliage.

TERIVATE

Apply 1-2 pts/A to burndown foliage prior to harvest. For best results, apply 1 pt/A followed by another 1 pt/A (minimum 5 day interval)

STREVIN 2EC

Apply Strevin 2EC at 3.2 – 5.8 oz/A to burndown foliage prior to harvest.

Tank-mixing – Apply Terivate at 1 pt/A and Strevin 2EC at 2.0 – 5.8 oz /A

- Devour®* (paraquat dichloride 3.0 lbs ai/gal) – **Fresh market potatoes only**
- For quick vine kill apply 1.3 pts/A
- For dense vines, apply 0.6 pts/A followed 5 days or more after with another 0.6 pts/A

* = Restricted Use Pesticide


ADJUVANT RECOMMENDATIONS

For best results use Aviator at 0.25% v/v or 1 qt/100 gal. This will provide spreading and penetration of the defoliant as well as superior water conditioning and acidification to enhance the burndown.


For added deposition and drift reduction use Adversary at 0.25% v/v or 1 qt/100 gal

Potato Vine Desiccation Trial

Twin Falls, Idaho 2023 | 3 Days after Treatment



Non-Treated



Diquat-Dibromide + Aviator

