

SP AZTEB™

Broad-spectrum fungicide for control of turf diseases on golf courses

ACTIVE INGREDIENTS	% BY WT
Azoxystrobin: methyl (E)-2-[[6-(2-cyanophenoxy)-4-pyrimidinyl]oxy]alpha-methoxymethylene benzeneacetate.....	11.9%
Tebuconazole: (+)-alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1 H-1,2,4-triazole-1-ethanol.....	26.9%
OTHER INGREDIENTS.....	61.2%
TOTAL.....	100.00%

A suspension concentrate fungicide containing 2.34 lb Tebuconazole and 1.04 lb Azoxystrobin per gal.

EPA REG. NO. 89442-27-95428 EPA EST. NO. 42750-MO-002

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional Precautionary Statements and Directions For Use inside booklet.
Reformulation is prohibited. See individual container labels for repackaging limitations.

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by the poison control center or doctor. • DO NOT give anything to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR ON CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center, or doctor, or when going for treatment. For emergency information concerning this product, call the poison control center at **1-800-222-1222**.

Manufactured For:
 Simplot Turf & Horticulture
 A Division of Simplot AB Retail, Inc.
 P.O. Box 9296, Boise, ID 83707
 (208) 336-2110

AVAILABLE IN: 1 GAL (3.79 L)

Guaranteed by:
 Simplot Turf & Horticulture
 P.O. Box 9296, Boise, ID 83707
 (208) 336-2110

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. HARMFUL IF SWALLOWED. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt and long pants, socks and shoes, and gloves. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemically resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separate from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing;
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Azoxystrobin can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

GROUND WATER ADVISORY

Azoxystrobin and tebuconazole are toxic to freshwater and estuarine/marine fish; and azoxystrobin is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

SURFACE WATER ADVISORY

This product may contaminate water through drift or spray in wind. This product has a high potential for runoff into surface waters for several days to weeks after application. Poorly draining soils, wet soils with readily visible slopes toward adjacent surface waters or soil with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours. Notify state and/or Federal authorities immediately if you observe any adverse environmental effects due to use of this product. Notify state and/or Federal authorities and Prime Source, LLC immediately if you observe any adverse environmental effects due to use of this product.

PHYSICAL HAZARDS

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. This label must be in the possession of the user at the time of pesticide application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Chemigation - Do not apply this product through any irrigation/chemigation system.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

PRODUCT INFORMATION

SP AzTeb™ is a combination of two broad spectrum, preventative fungicides with systemic properties for the control of many important turfgrass diseases on golf course turf only. SP AzTeb™ may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered, turf protection products. All applications must be made according to the use directions that follow. This product is not for residential use.

PRODUCT TANK MIX USE PRECAUTIONS

SP AzTeb™ has demonstrated some phytotoxic effects when mixed with products that are formulated as EC's. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

INTEGRATED PEST (DISEASE) MANAGEMENT

Integrate SP AzTeb™ into an overall disease and pest management strategy whenever the use of a fungicide is required. Follow cultural practices known to reduce disease development. The USE DIRECTIONS section in this label identifies specific IPM instructions. Consult your local turf authority for additional IPM strategies established for your area. SP AzTeb™ may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

RESISTANCE MANAGEMENT

- SP AzTeb™ is a mixture of a Group 3 (tebuconazole) and a Group 11 (azoxystrobin) fungicide. SP AzTeb™ has two modes of action: (1) DMI (Demethylation Inhibitor of sterol biosynthesis) [Group 3], and (2) inhibitor of the Qo (quinone outside) site within the electron transport system (QoI) as well as disrupting membrane synthesis by blocking demethylation [Group 11]. Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. SP AzTeb™ is not cross-resistant with other classes of fungicides which have different modes of action. Because resistance development cannot be predicted, use of this product must conform to resistance management strategies established for turf in your area. Consult your local or State turf authorities for resistance management strategies that are complementary to those in this label.
- Resistance management strategies may include rotating and/or tank-mixing with products having different modes of action or limiting the total number of applications per season. Avoid alternation with other QoIs, such as pyraclostrobin and trifloxystrobin or other DMI's such as triadimefon, fenarimol, and myclobutanil. Do not alternate or tank mix SP AzTeb™ with fungicides to which resistance has already developed. Continual use of SP AzTeb™ may allow less sensitive strains of pathogens to increase in the population and reduce the efficacy of SP AzTeb™. Prime Source, LLC encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

- Use of a multiple-site contact fungicide such as Daconil as an alternation or tank-mix partner can help alleviate selection pressure on pathogens and delay the onset of resistance.
- Since pathogens differ in their potential to develop resistance to SP AzTeb™, refer to the **USE DIRECTIONS** section in this label provides resistance management strategies specific for each disease.

SPRAYING/MIXING

- Do not apply SP AzTeb™ to turf by air.
- Apply SP AzTeb™ with all types of spray equipment commonly used for making ground applications. Do not apply SP AzTeb™ through any type of ultra low volume (ULV) spray system (less than 3 gal per acre). The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when disease conducive environmental conditions exist.
- Apply SP AzTeb™ in sufficient water volume for adequate coverage.

SPRAY DRIFT MANAGEMENT

- **Bermudagrass can be sensitive to SP AzTeb™ .**
- SP AzTeb™ is extremely phytotoxic to certain apple varieties. AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).
- Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

RESTRICTIONS

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

- DO NOT apply by ground within 100 ft of aquatic areas listed above.
- DO NOT cultivate within 10 ft of an aquatic area to allow growth of a vegetative filter strip.
- DO NOT spray SP AzTeb™ where spray drift may reach apple trees.
- DO NOT apply when weather conditions favor drift from treated areas to non-target aquatic habitat.
- DO NOT spray when conditions favor drift beyond area intended for application.
- DO NOT use spray equipment which has been previously used to apply SP AzTeb™ to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.
- DO NOT apply SP AzTeb™ to turf by air.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

APPLICATION INSTRUCTIONS

Apply SP AzTeb™ at rates and timings as described in this label.

MIXING INSTRUCTIONS

- To prepare spray solution, partially fill the spray tank with clean water and begin agitation. Add the specified amount of SP AzTeb™ to the tank, allowing time for good dispersion, then add an adjuvant, if recommended. If tank mixes are required, product should be added to the spray tank in the following order; SP AzTeb™, wettable granular or dry flowable formulations, wettable powders and flowable (aqueous suspensions) products. Finish filling the tank to the desired volume to obtain the proper spray concentration. Maintain agitation throughout the spraying operation.
- Do not allow spray mixture to stand overnight or for prolonged periods. Make up only the amount of spray required for immediate use. Sprayers should be thoroughly cleaned immediately after application.
- Do not use silicone-based products with SP AzTeb™ due to possible phytotoxicity.
- SP AzTeb™ is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides and biological control products, if tank mixes are desired, observe all directions, precautions* and limitations on labeling of all products used.
- Consult compatibility charts or your local or State turf authority for compatibility information. Do not combine SP AzTeb™ in the spray tank with pesticides, surfactants, or fertilizers, unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective, and non-injurious under your conditions of use. If physical compatibility is unknown, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least 20 minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

USE DIRECTIONS

SP AzTeb™ is for control of certain pathogens causing foliar, stem, and root diseases, including leaf and stem blights, leaf spots, patch diseases, mildews, anthracnose, fairy rings, molds, and rusts of turfgrass in golf courses. This product is not for residential use. Do not apply this product by aerial application.

Integrated Pest (Disease) Management (IPM):

Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management should be integrated with the use of fungicides to increase turf vigor and reduce the susceptibility to disease. Immunoassay detection kits and extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

Application Directions:

- Apply SP AzTeb™ prior to disease development. Mix SP AzTeb™ with the required amount of water and apply as a dilute spray application in 2 - 4 gal of water per 1,000 sq ft (87-174 gal per acre).
- Repeat applications at specified intervals for as long as required, not to exceed the maximum specified use rate per season. For spot treatments, use 0.75 fl oz SP AzTeb™ per 1 to 2 gal of water.

For Use with Soil Injection Applications:

Apply SP AzTeb™ through a liquid fungicide injector for the control of ectotrophic root diseases such as summer patch and take-all patch. Use SP AzTeb™ only in liquid injection equipment specifically designated for pesticide use.

For Use with Soil Injection Applications:

Apply SP AzTeb™ at 0.75 - 1.7 fl oz per 1,000 sq ft. Spray carrier volume should fall within 30-150 gal of water per 1,000 sq ft. Injection hole spacing of 1-in by 1-in is required for optimum control. Injection depth must not be greater than 2-in. One inch depth is required for optimum results. Application should follow disease control strategies used for normal broadcast spray programs.

For Use in the Establishment of Turfgrass from Seed or in Overseeding of Dormant Turfgrass:

Apply SP AzTeb™ to control of certain turfgrass diseases associated with turfgrass establishment from seed. SP AzTeb™ may also be used during overseeding of dormant turfgrass.

Apply SP AzTeb™ before or after seeding or at seedling germination and emergence to ryegrass, bentgrass, bluegrass, and fescue turfgrass types. Optimum application timing is during seeding. See Application Directions section.

Rate Ranges:

Use the shorter specified application interval and/or use the higher specified rate when prolonged favorable disease conditions exist.

TURF USE RESTRICTIONS

- **For use on golf course turf only.**
- Not for residential use.
- Not for use on turf being grown for sale or commercial use as sod.
- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
- Do not graze or feed clippings from treated turf areas to animals.
- Do not apply SP AzTeb™ to turf by air. Applications must be made by ground only.
- Do not exceed 1.5 fl oz per 1,000 sq. ft. every 30 days on any variety of bermudagrass.
- In FL, do not apply SP AzTeb™ to bermudagrass golf course greens when temperatures exceed 90 °F.
- Do not apply more than 1.75 gal of product per acre per year (5.1 fl oz per 1,000 sq ft per year).
- Do not apply more than 6 applications per year, except in NY State.
- Do not apply more than 3 applications at 0.75 fl oz of SP AzTeb™ per 1,000 sq ft per year in New York State. Do not exceed a total of 98 fl oz (1.8 lb of tebuconazole A.I.) per acre per year in New York State.

TARGET DISEASES	USE RATE FL OZ PRODUCT PER 1000 SQ FT	APPLICATION INTERVAL (DAYS)	REMARKS*
Anthracnose (<i>Colletotrichum cereale</i>) (formerly known as <i>C. graminicola</i>)	0.75 - 1.5	14 - 28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development. The maximum application interval for greens is 14 days.
Bentgrass Dead Spot (<i>Ophiosphaerella agrostis</i>)	0.75 - 1.5	14	Begin applications prior to disease development.
Brown Patch (<i>Rhizoctonia solani</i>)	0.5 - 0.75	14	For higher cut cool season turfgrass, St. Augustine grass and centipedegrass (above 0.375 in. in height). Apply when conditions are favorable for disease development.
Brown Patch (<i>Rhizoctonia solani</i>) and other Rhizoctonia (<i>R. spp.</i>) diseases such as Large Patch, Zoysia Patch, Leaf & Sheath Spot (<i>R. zeae</i>)	0.75 - 1.5	14 - 28	For Brown Patch, apply when conditions are favorable for disease development. For Large Patch of all warm season turfgrasses, make 1 or 2 applications in the fall prior to infection or when conditions are favorable for infection. For Zoysia Patch, make 1 or 2 applications approximately one month prior to zoysiagrass dormancy. Reapply 14 to 28 days later. For Leaf & Sheath Spot (<i>R. zeae</i>), apply when conditions are favorable for infection such as sequential or periods of temperatures at or above 90 degrees Fahrenheit. Curative control may necessitate several applications. Target spray applications at crown of turfgrass.
Cool Weather Brown Patch, Yellow Patch (<i>Rhizoctonia cerealis</i>)	1.5	28	Make 1 or 2 applications in fall or when conditions are favorable for disease development.
Dichondra Rust (<i>Puccinia dichondrae</i>)	1.5	14 - 21	Apply when conditions are favorable for disease development.
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	0.75 - 1.5	7 - 14	Apply preventively on higher cut cool season turfgrass (fairways, lawns, etc.) mowed at or above 0.375 in. in height during periods of low to moderate disease pressure.
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	0.75 - 1.5	14 - 28	Apply preventively when conditions are favorable for disease development. For turf with active dollar spot, use the higher rate and combine with Daconil.
Fairy Ring (<i>Lycoperdon spp.</i> , <i>Agrocybe pediades</i> , and <i>Bovista plumbea</i>)	0.75 - 1.5	14 - 28	Apply as soon as possible after fairy ring symptoms develop. Apply in 2-4 gal of water per 1,000 sq ft. Add the specified rate of a wetting agent to the final spray. Severely damaged or thin turf may require reseeding. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. If area is hydrophobic, use wetting agents and irrigate prior to application(s) of SP AzTeb™. Reapplication after 28 days may be required in some cases.
Gray Leaf Spot (<i>Pyricularia grisea</i>)	0.75 - 1.5	14 - 28	Use SP AzTeb™ in a preventive disease control program. Begin applications before disease is present and alternate with other fungicide chemistries that control gray leaf spot.
Gray Snow Mold, Typhula Blight (<i>Typhula incarnata</i>)	1.5	10 - 28	Make two applications of 1.5 fl oz spaced 10-28 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide, such as Daconil and Medallion, may enhance control under severe disease pressure.
Leaf Rust, Stem Rust, Stripe Rust (<i>Puccinia spp.</i>)	0.75 - 1.5	14 - 28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Leaf Spot (<i>Bipolaris sorokiniana</i>)	0.75 - 1.5	14 - 21	Apply when conditions are favorable for disease development.
Melting Out (<i>Drechslera poae</i>)	0.75 - 1.5	14 - 21	Apply when conditions are favorable for disease development.
Microdochium Patch (formerly known as Fusarium Patch) (<i>Microdochium nivale</i>)	0.75 - 1.5	14 - 28	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Necrotic Ring Spot (<i>Ophiosphaerella korrae</i>)	1.5	14 - 28	Apply when conditions are favorable for disease development.
Pink Patch (<i>Limonomyces roseipellis</i>)	0.75 - 1.5	14 - 28	Apply when conditions are favorable for disease development.

TARGET DISEASES	USE RATE FL OZ PRODUCT PER 1000 SQ FT	APPLICATION INTERVAL (DAYS)	REMARKS*
Pink Snow Mold (<i>Microdochium nivale</i>)	1.5	10 - 28	Make two applications of 1.5 fl oz spaced 10-28 days apart in late fall just before snow cover. Tank mixing with another snow mold fungicide, such as Daconil and Medallion, may enhance control under severe disease pressure.
Powdery Mildew (<i>Erysiphe graminis</i>)	0.75 - 1.5	14 - 28	Begin applications when conditions are favorable for disease infection, prior to disease symptom development.
Pythium Blight, Pythium Root Rot (<i>Pythium aphanidermatum</i> , <i>Pythium spp.</i>)	1.5	10 - 14	Use preventatively. Begin applications when conditions are favorable for disease infection prior to disease symptom development. During periods of prolonged favorable conditions, treat on the 10-day application interval. For use on newly seeded well as established turf.
Red Thread (<i>Laetisaria fuciformis</i>)	0.75 - 1.5	14 - 28	Apply when conditions are favorable for disease development.
Southern Blight (<i>Sclerotium rolfsii</i>)	0.75 - 1.5	14 - 28	Apply when conditions are favorable for disease development.
Spring Dead Spot (<i>Ophiosphaerella korrae</i>) or (<i>Ophiosphaerella narmari</i>) or (<i>Ophiosphaerella herpotricha</i>)	1.5	14 - 28	Apply 1 or 2 applications approximately one month prior to bermudagrass dormancy. ¼ in. to ½ in. of irrigation directly after application is recommended. Reapply 14 to 28 days later.
Summer Patch (<i>Magnaporthe poae</i>)	0.75 - 1.5	14 - 28	Apply when conditions are favorable for disease development. Initiate applications when soil temperatures reach 65 °F at a 2 in. soil depth.
Take-all Patch (<i>Gaeumannomyces graminis var. venae</i>)	1.5	28	Begin applications when conditions are favorable for disease infection prior to disease symptom development. Make two applications, 28 days apart in the spring and two applications 28 days apart in the fall.

*Do not apply more than two sequential applications of SP AzTeb™ for control of Gray Leaf Spot and *Pythium spp.* For all other diseases when Gray Leaf Spot and *Pythium spp.* are not present, do not apply more than three sequential applications of SP AzTeb™.

SP AzTeb™ RATE CONVERSION CHART FOR TURF

FL OZ PRODUCT* PER 1,000 SQ FT	PINTS PRODUCT PER ACRE	GAL PRODUCT PER ACRE
0.75	2.0	0.26
1.0	2.7	0.34
1.5	4.0	0.51

*One (1) fl oz of SP AzTeb™ contains 0.2925 oz (weight) of tebuconazole and 0.1300 oz (weight) azoxystrobin.

AMOUNT OF SP AzTeb™ FOR TURF APPLICATIONS PER 100 GAL

SPRAY VOLUME: 2 GAL PER 1,000 SQ FT

SP AzTeb™ USE RATE (FL OZ PER 1,000 SQ FT)	PINTS OF PRODUCT PER 100 GAL
0.75	2.3
1.0	3.1
1.5	4.7

SPRAY VOLUME: 3 GAL PER 1,000 SQ FT

SP AzTeb™ USE RATE (FL OZ PER 1,000 SQ FT)	PINTS OF PRODUCT PER 100 GAL
0.75	1.5
1.0	2.1
1.5	3.0

SPRAY VOLUME: 4 GAL PER 1,000 SQ FT

SP AzTeb™ USE RATE (FL OZ PER 1,000 SQ FT)	PINTS OF PRODUCT PER 100 GAL
0.75	1.15
1.0	1.55
1.5	2.25

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop up and remove to chemical waste storage area until proper disposal can be made if product cannot be reused according to the label.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance. Open dumping is prohibited.

CONTAINER HANDLING [less than 5 gallons]: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Handling [greater than 5 gallons]: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IN CASE OF EMERGENCY CALL CHEMTREC 1-800-424-9300

© 2025 J.R. Simplot Company. All rights reserved. SIMPLOT[®] and SP AzTeb[™] are trademarks in the United States and other countries and owned by the J.R. Simplot Company. 25196 530041G F3237

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties, and Limitations of Liability. **CONDITIONS, DISCLAIMER OF WARRANTIES, and LIMITATIONS OF LIABILITY. CONDITIONS:** The Directions For Use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Simplot AB Retail, Inc. and its affiliates (collectively, "Simplot"). To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Simplot makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Simplot is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Simplot disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or, at Simplot's election, the replacement of product.